

20 . Pathogen Genomics Center

- 1) Ode H, Yokoyama M, Kanda T, and Sato H. Identification of folding preferences of cleavage junctions of HIV-1 precursor proteins for regulation of cleavability. *J. Mol. Model.* In press (2010)
- 2) Onyango CO, Leligdowicz A, Yokoyama M, Sato H, Song H, Nakayama EE, Shioda T, de Silva T, Townend J, Jaye A, Whittle H, Rowland-Jones S, Cotten M. HIV-2 capsids distinguish high and low virus load patients in a West African community cohort. *Vaccine.* 28 Suppl 2:B60-7, 2010.
- 3) Gatanaga H, Ode H, Hachiya A, Hayashida T, Sato H, Takiguchi M, Oka S. Impact of human leukocyte antigen-B*51-restricted cytotoxic T-lymphocyte pressure on mutation patterns of nonnucleoside reverse transcriptase inhibitor resistance. *AIDS.* 24(5):F15-22, 2010.
- 4) Gatanaga H, Ode H, Hachiya A, Hayashida T, Sato H, Oka S. Combination of V106I and V179D polymorphic mutations in human immunodeficiency virus type 1 reverse transcriptase confers resistance to efavirenz and nevirapine but not etravirine. *Antimicrob Agents Chemother.* 54:1596-602, 2010.
- 5) Yokoyama M, Mori H, Sato H. Allosteric regulation of HIV-1 reverse transcriptase by ATP for nucleotide selection.. *PLoS One.* 5:e8867, 2010.
- 6) Nakajima N, Hata S, Sato Y, Tobiume M, Katano H, Kaneko K, Nagata N, Kataoka M, Aina A, Hasegawa H, Tashiro M, Kuroda M, Odai T, Urasawa N, Ogino T, Hanaoka H, Watanabe M, Sata T. The first autopsy case of pandemic influenza (A/H1N1pdm) virus infection in Japan: detection of a high copy number of the virus in type II alveolar epithelial cells by pathological and virological examination. *Jpn J Infect Dis.* 63:67-71, 2010.
- 7) Okutani A, Sekizuka T, Boldbaatar B, Yamada A, Kuroda M, Inoue S. Phylogenetic typing of *Bacillus anthracis* isolated in Japan by multiple locus variable-number tandem repeats and the comprehensive single nucleotide polymorphism. *J Vet Med Sci.* 72: 93-97. 2010.
- 8) Sato, H., Kusumoto-Matsuo, R., Ishii, Y., Mori, S., Nakahara, T., Shinkai-Ouchi, F., Kawana, K., Fujii, T., Taketani, Y., Kanda, T. and Kukimoto, I. Identification of nucleolin as a protein that binds to human papillomavirus type 16 DNA. *Biochem Biophys Res Commun,* 387, 525-530, 2009.
- 9) Kondo, K, Ishii, Y, Mori, S, Shimabukuro, S, Yoshikawa, H, Kanda, T. Nuclear location of minor capsid protein L2 is required for expression of a reporter plasmid packaged in HPV51 pseudovirions. *Virology,* 394, 259-65, 2009.
- 10) McCord, RA., Michishita, E., Hong, T., Berber, E., Boxer, LD., Kusumoto, R., Guan, S., Shi, X., Gozani, O., Burlingame, AL., Bohr, VA. and Chua, KF. SIRT6 stabilizes DNA-dependent Protein Kinase at chromatin for DNA double-strand break repair. *Aging (Albany NY),* 1, 109-121, 2009.
- 11) Oka T, Yokoyama M, Katayama K, Tsunemitsu H, Yamamoto M, Miyashita K, Ogawa S, Motomura K, Mori H, Nakamura H, Wakita T, Takeda N, Sato H. Structural and biological constraints on diversity of regions immediately upstream of cleavage sites in calicivirus precursor proteins. *Virology.* 394:119-29. 2009.
- 12) Sahbandar IN, Takahashi K, Djoerban Z, Firmansyah I, Naganawa S, Motomura K, Sato H, Kitamura K, Pohan HT, Sato S. Current HIV type 1 molecular epidemiology profile and identification of unique recombinant forms in Jakarta, Indonesia. *AIDS Res Hum Retroviruses.* 25:637-46, 2009.
- 13) Kamiyama H, Yoshii H, Tanaka Y, Sato H, Yamamoto N, Kubo Y. Raft localization of CXCR4 is primarily required for X4-tropic human immunodeficiency virus type 1 infection. *Virology* 386:23-31, 2009.

