

## 11 . Department of Medical Entomology

- 1)Sasaki T. Hiraoka T. Kobayashi M. Hemolytic activity is mediated by the endogenous lectin in the mosquito hemolymph serum. *J. Insect Physiol.* 56(9): 1032-1039, 2010.
- 2) Komagata, O., Kasai, S., Tomita, T. Overexpression of cytochrome P450 genes in pyrethroid-resistant *Culex quinquefasciatus*. *Insect Biochem. Mol. Biol.* 40:146-152, 2010.
- 3)Itokawa, K., Komagata, O., Kasai, S., Okamura, Y., Masada, M., Tomita, T. Genomic structures of *Cyp9m10* in pyrethroid resistant and susceptible strains of *Culex quinquefasciatus*. *Insect Biochem. Mol. Biol.* 40: 631-640, 2010.
- 4)Tsuda, Y. and Kim, K.S. Prediapause migration and overwintering of *Culex tritaeniorhynchus* (Diptera: Culicidae) observed in a park in urban Tokyo during 2007 and 2009. *Med. Entomol. Zool.* 61: 69-78 ,2010.
- 5)Hoshino K., Isawa, H., Tsuda Y. and Kobayashi M. Laboratory Colonization of *Aedes japonicus japonicus* (Diptera: Culicidae) collected in Narita, Japan and biological properties of the established colony. *Jpn. J. Infect. Dis.* 63: 401-404 ,2010.
- 6)Kim, K.S. and Tsuda Y. Seasonal changes in feeding pattern of *Culex pipiens pallens* govern transmission dynamics of multiple lineages of avian malaria parasite in Japanese wild bird community. *Mol. Ecol.* 19: 5545-5554,2010.
- 7)Higa, Y., Toma, T., Tsuda, Y. and Miyagi, I. A multiplex PCR-based molecular identification of five morphologically related, medically important subgenus *Stegomyia* mosquitoes from the Genus *Aedes* (Diptera: Culicidae) found in the Ryukyu Archipelago, Japan. *Jpn. J. Infect. Dis.* 63: 312-316,2010.
- 8)Somboon, P., Rory, A., Tsuda, Y., Takagi, M. and Harbach, R. E. Systematics of *Anopheles* (*Cellia*) *yaeyamaensis* sp. n., alias species E of the *An. minimus* complex in southeastern Asia (Diptera: Culicidae). *Zootaxa* 2651: 43–51 ,2010.
- 9)Yotsu, R. R., Tamaki, T., Ujiie, M., Takeshita, N., Kanagawa, S., Kudo, K. and Hayashi, T. Imported tungiasis in a Japanese student returning from East Africa. *J. Dermatol.* no. doi : 10.1111 /j.1346-8138.2010.00952.x,2010.
- 10)Hayashi, T. The genus *Opalimosina* Roháček (Diptera, Sphaeroceridae) from Japan. *Med. Entomol. Zool.* 61: 309-319, 2010.
- 11)Kuwata R., Isawa H., Hoshino K., Tsuda Y., Yanase T., Sasaki T., Kobayashi M., Sawabe K. RNA splicing in a new Rhabdovirus from *Culex* mosquitoes. *J. Virol.* 85: 6185-6196, 2011.
- 12)Isawa H., Kuwata R., Hoshino K., Tsuda Y., Sakai K., Watanabe S., Nishimura M., Satho T., Kataoka M., Nagata N., Hasegawa H., Bando H., Yano K., Sasaki T., Kobayashi M., Mizutani T. Sawabe K. Identification and molecular characterization of a new nonsegmented double-stranded RNA virus isolated from *Culex* mosquitoes in Japan. *Virus Res.* 155: 147-155, 2011.
- 13)Sawabe K., Hoshino K., Isawa H., Sasaki T., Kim K.S., Hayashi T., Tsuda Y., Kurahashi H. Kobayashi M. Blow flies were one of the possible candidates for transmission of highly pathogenic H5N1 avian influenza virus during the 2004 outbreaks in Japan. *Influenza Res. Treat.* 2011: 652652, 2011.
- 14)Sawabe K., Isawa H., Hoshino K., Sasaki T., Roychoudhury S., Higa Y., Kasai S., Tsuda Y., Nishiumi I., Hisai N., Hamao S. Kobayashi M. Host-feeding habits of *Culex pipiens* and *Aedes albopictus* (Diptera: Culicidae) collected at the urban and suburban residential areas of Japan. *J. Med. Entomol.* 47: 442-450, 2010.
- 15)Imura T., Sato Y., Ejiri H., Tamada A., Isawa H., Sawabe K., Omori S., Murata K. Yukawa M. Molecular identification of blood source animals from black flies (Diptera: Simuliidae) collected in the alpine regions of Japan. *Parasitol. Res.* 106: 543-547, 2010.
- 16)Hardstone, M.C., Komagata, O., Kasai, S., Tomita, T., Scott, J.G. Use of isogenic strains indicates CYP9M10 is linked to permethrin resistance in *Culex pipiens quinquefasciatus*. *Insect Mol. Biol.* 19: 717-726, 2010.