

## 1. Department of Virology I

- 1) Saijo M. Dual use research of concern issues in the field of Microbiology Research in Japan. *Journal of Disaster Research*. 2013;8:693-7.
- 2) Yamanaka A, Iwakiri A, Yoshikawa T, Sakai K, Singh H, Himeji D, Kikuchi I, Ueda A, Yamamoto S, Miura M, Shioyama Y, Kawano K, Nagaishi T, Saito M, Minomo M, Iwamoto N, Hidaka Y, Sohma H, Kobayashi T, Kanai Y, Kawagishi T, Nagata N, Fukushi S, Mizutani T, Tani H, Taniguchi S, Fukuma A, Shimojima M, Kurane I, Kageyama T, Odagiri T, Saijo M, Morikawa S. Imported case of acute respiratory tract infection associated with a member of species nelson bay orthoreovirus. *PLoS One*. 2014;9(3):e92777.
- 3) Shimojima M, Takenouchi A, Shimoda H, Kimura N, Maeda K. Distinct usage of three C-type lectins by Japanese encephalitis virus: DC-SIGN, DC-SIGNR, and LSECtin. *Arch Virol*. 2014;159(8):2023-31.
- 4) Sakai K, Nagata N, Ami Y, Seki F, Suzaki Y, Iwata-Yoshikawa N, Suzuki T, Fukushi S, Mizutani T, Yoshikawa T, Otsuki N, Kurane I, Komase K, Yamaguchi R, Hasegawa H, Saijo M, Takeda M, Morikawa S. Lethal canine distemper virus outbreak in cynomolgus monkeys in Japan in 2008. *J Virol*. 2013;87(2):1105-14.
- 5) Sakai K, Yoshikawa T, Seki F, Fukushi S, Tahara M, Nagata N, Ami Y, Mizutani T, Kurane I, Yamaguchi R, Hasegawa H, Saijo M, Komase K, Morikawa S, Takeda M. Canine distemper virus associated with a lethal outbreak in monkeys can readily adapt to use human receptors. *J Virol*. 2013;87(12):7170-5.
- 6) Takahashi T, Maeda K, Suzuki T, Ishido A, Shigeoka T, Tominaga T, Kamei T, Honda M, Ninomiya D, Sakai T, Senba T, Kaneyuki S, Sakaguchi S, Satoh A, Hosokawa T, Kawabe Y, Kurihara S, Izumikawa K, Kohno S, Azuma T, Suemori K, Yasukawa M, Mizutani T, Omatsu T, Katayama Y, Miyahara M, Ijuin M, Doi K, Okuda M, Umeki K, Saito T, Fukushima K, Nakajima K, Yoshikawa T, Tani H, Fukushi S, Fukuma A, Ogata M, Shimojima M, Nakajima N, Nagata N, Katano H, Fukumoto H, Sato Y, Hasegawa H, Yamagishi T, Oishi K, Kurane I, Morikawa S, Saijo M. The first identification and retrospective study of severe fever with thrombocytopenia syndrome in Japan. *J Infect Dis*. 2014;209(6):816-27.
- 7) Moi ML, Omatsu T, Hirayama T, Takasaki T, Nakamura S, Tajima S, Ito M, Yoshida T, Saito A, Katakai Y, Akari H, Kurane I. Presence of viral genome in urine and development of hematuria in common marmosets (*Callithrix jacchus*) after inoculation with dengue virus. *Pathogens*. 2013;2:357-63.
- 8) Moi ML, Takasaki T, Kurane I. Efficacy of tetravalent dengue vaccine in Thai schoolchildren. *Lancet*. 2013;381,9872,1094. <Letter to Editor>
- 9) Moi ML, Omatsu T, Tajima S, Lim CK, Kotaki A, Ikeda M, Harada F, Ito M, Saijo M, Kurane I, Takasaki T. Detection of dengue virus non-structural protein 1 (NS1) by using ELISA as a useful laboratory diagnostic method for dengue virus infection of international travellers. *J Travel Med*. 2013;20,3:185-93.
- 10) Moi ML, Lim CK, Takasaki T, Kurane I. Infection-enhancement activity in dengue patients using undiluted serum samples. *Trans R Soc Trop Med Hyg*. 2013;107:5-58.
- 11) Tochitani K, Shimizu T, Shinohara K, Tsuchido Y, Moi ML, Takasaki T. Ross River virus - Japan ex Australia: (VI). ProMed, promed archive no. 2013;20130616;1776324.
- 12) Mizuno Y, Takasaki T. ProMed-mail posts. Dengue case\_Japan ex Kenya (Mombasa) . *Dengue/DHF update*. 2013;(58):2.
- 13) Takeshita N, Lim CK, Mizuno Y, Shimbo T, Kotaki A, Ujiie M, Hayakawa K, Kato Y, Kanagawa S, Kaku M, Takasaki T. Immunogenicity of single-dose Vero cell-derived Japanese encephalitis vaccine in Japanese adults. *J Infect Chemother*. 2014;20:238-42.
- 14) Trent DW, Minor P, Jivapaisarnpong T, Shin J. WHO working group on the quality, safety and efficacy of Japanese encephalitis vaccines (live attenuated) for human use, Bangkok, Thailand, 21e23 February 2012. *Biologicals* 2013;41; 450-7.

- 15) Suzuki R, Ishikawa T, Konishi E, Matsuda M, Watashi K, Aizaki H, Takasaki T, Wakita T. Production of single-round infectious chimeric flaviviruses with DNA-based Japanese encephalitis virus replicon. *J Gen Virol.* 2013;95:60-5.
- 16) Nidaira M, Kyan H, Taira K, Okano S, Oshiro T, Kato T, Kudo N, Azama Y, Mahoe Y, Kudaka J, Tamanaha K, Takasaki T. Survey of Japanese encephalitis virus in pigs and wild boars on Ishigaki and Iriomote Islands in Okinawa, Japan. *Epidemiol Infect.* 2013;5:1-5.
- 17) Kurane I, Shibasaki K, Kotaki A, Hijioka Y, Takasaki T. The Effect of Precipitation on the Transmission of Japanese Encephalitis (JE) Virus in Nature: A Complex Effect on Antibody-Positive Rate to JE Virus in Sentinel Pigs. *Int J Environ Res Public Health* 2013. 2013;10:1831-44.
- 18) Noyori O, Nakayama E, Maruyama J, Yoshida R, Takada A. Suppression of Fas-mediated apoptosis via steric shielding by filovirus glycoproteins. *Biochem Biophys Res Commun.* 2013;441(4):994-8.
- 19) Noyori O, Matsuno K, Kajihara M, Nakayama E, Igarashi M, Kuroda M, Isoda N, Yoshida R, Takada A. Differential potential for envelope glycoprotein-mediated steric shielding of host cell surface proteins among filoviruses. *Virology.* 2013;446(1-2):152-61.
- 20) Nakayama E, Saijo M. Animal models for Ebola and Marburg virus infections. *Front Microbiol.* 2013;4:267.
- 21) Kajihara M, Nakayama E, Marzi A, Igarashi M, Feldmann H, Takada A. Novel mutations in Marburg virus glycoprotein associated with viral evasion from antibody mediated immune pressure. *J Gen Virol.* 2013;94(Pt 4):876-83.
- 22) Moi ML, Takasaki T, Saijo M, Kurane I. Determination of antibody concentration as the main parameter in a dengue virus antibody-dependent enhancement assay using FcγR2A-expressing BHK cells. *Arch Virol.* 2014;159(1):103-16.
- 23) Moi ML, Takasaki T, Omatsu T, Nakamura S, Katakai Y, Ami Y, Suzaki Y, Saijo M, Akari H, Kurane I. Demonstration of marmosets (*Callithrix jacchus*) as a non-human primate model for secondary dengue virus infection: high levels of viremia and serotype cross-reactive antibody responses consistent with secondary infection of humans. *J Gen Virol.* 2014;42(1):591-600.
- 24) Takayama-Ito M, Nakamichi K, Kinoshita H, Kakiuchi S, Kurane I, Saijo M, Lim CK. Sensitive in vitro assay for the detection of residual viable rabies virus in inactivated rabies vaccines. *Biologicals* 2014;42:42-7.
- 25) Nakamichi K, Inoue N, Shimokawa T, Kurane I, Lim CK, Saijo M. Detection of human herpesviruses in the cerebrospinal fluid from patients diagnosed with or suspected of having progressive multifocal leukoencephalopathy. *BMC Neurology* 2013;13:200 (online).
- 26) Mikita K, Maeda T, Fujikura Y, Kozaki Y, Hara Y, Kanoh S, Kishida S, Saijo M, Nakamichi K, Kawana A. Does anti-JCV therapy improve the prognosis of AIDS-related PML? *Clin Neurol Neurosurg.* 2013;115(9):1853-4.
- 27) Nagayama S, Gondo Y, Araya S, Minato N, Fujita-Nakata M, Kaito M, Nakanishi M, Tanaka K, Yamaya H, Yokoyama H, Nakamichi K, Saijo M, Okamoto K, Toyoshima Y, Kakita A, Matsui M. Progressive multifocal leukoencephalopathy developed 26 years after renal transplantation. *Clin Neurol Neurosurg.* 2013;115(8):1482-4.
- 28) Takeshita N, Lim CK, Mizuno Y, Shimbo T, Kotaki A, Ujiie M, Hayakawa K, Kato Y, Kanagawa S, Kaku M, Takasaki T. Immunogenicity of single-dose Vero cell-derived Japanese encephalitis vaccine in Japanese adults. *J Infect Chemother.* 2014;20(238-42).
- 29) Nakamichi K, Tajima S, Lim CK, Saijo M. High-resolution melting analysis for mutation scanning in the non-coding control region of JC polyomavirus from patients with progressive multifocal leukoencephalopathy. *Arch Virol.* 2014;159:1687-96.
- 30) Wang LX, Takayama-Ito M, Kinoshita-Yamaguchi H, Kakiuchi S, Suzutani T, Nakamichi K, Lim CK, Kurane, Saijo M. Characterization of DNA polymerase-associated acyclovir-resistant herpes simplex virus type 1: mutations,

- sensitivity to antiviral compounds, neurovirulence, and in-vivo sensitivity to treatment. *Jpn J Infect Dis.* 2013;66(5):404-10.
- 31) Moi ML, Omatsu T, Tajima S, Lim CK, Kotaki A, Ikeda M, Harada F, Ito M, Saijo M, Kurane I, Takasaki T. Detection of dengue virus nonstructural protein 1 (NS1) by using ELISA as a useful laboratory diagnostic method for dengue virus infection of international travelers. *J Travel Med.* 2013;20(3):185-93.
- 32) Kakiuchi, S, Nonoyama S, Wakamatsu H, Kogawa K, Wang, L, Kinoshita-Yamaguchi H, Takayama-Ito M, Lim CK, Inoue N, Mizuguchi M, Igarashi T, Saijo M. Neonatal herpes encephalitis caused by a virologically confirmed acyclovir-resistant herpes simplex virus 1 strain. *J Clin Microbiol.* 2013;51(1):356-9.
- 33) Nakamichi K, Kishida S, Tanaka K, Suganuma A, Sano Y, Sano H, Kanda T, Maeda N, Kira J, Itoh A, Kato N, Tomimoto H, Kurane I, Lim C.K, Mizusawa H, Saijo M. Sequential changes in the non-coding control region sequences of JC polyomaviruses from the cerebrospinal fluid of patients with progressive multifocal leukoencephalopathy. *Arch Virol.* 2013;158(3):639-50.
- 34) Hashimoto K, Yamada S, Katano H, Fukuchi S, Sato Y, Kato M, Yamaguchi T, Moriishi K, Inoue N. Effects of immunization of pregnant guinea pigs with guinea pig cytomegalovirus glycoprotein B on viral spread in the placenta. *Vaccine.* 2013;31:3199-3205.
- 35) Ikuta K, Minematsu T, Inoue N, Kubo T, Asano K, Ishibashi K, Imamura T, Nakai H, Yoshikawa T, Moriuchi H, Fujiwara S, Koyano S, Suzutani T. Cytomegalovirus (CMV) glycoprotein H-based serological analysis in Japanese healthy pregnant women, and in neonates with congenital CMV infection and their mothers. *J Clin Virol.* 2013;58:474-8.
- 36) Nakamura H, Liao H, Minami K, Toyoda M, Akutsu H, Miyagawa Y, Okita H, Kiyokawa N, Umezawa A, Imadome K, Inoue N, Fujiwara S. Human cytomegalovirus induces apoptosis in neural stem/progenitor cells derived from induced pluripotent stem cells by generating mitochondrial dysfunction and endoplasmic reticulum stress. *Herpesviridae.* 2013;4:2.
- 37) Taniguchi R, Koyano S, Suzutani T, Goishi K, Ito Y, Morioka I, Oka A, Nakamura H, Yamada H, Igarashi T, Inoue N. Polymorphisms in Toll-like receptor 2 are associated with congenital cytomegalovirus infection. *Int J Infect Dis.* 2013;17:e1092-7.
- 38) Andoh M, Andoh R, Teramoto K, Komiya T, Kaneshima T, Takano A, Hayashidani H, Ando S. Survey of *Coxiella burnetii* in ticks collected from dogs in Japan. *J Vet Med Sci.* 2013;75(8):1115-7.
- 39) Kawabata H, Takano A, Kadosaka T, Fujita H, Nitta Y, Gokuden M, Honda T, Tomida J, Kawamura Y, Masuzawa T, Ishiguro F, Takada N, Yano Y, Andoh M, Ando S, Sato K, Takahashi H, Ohnishi M. Multilocus sequence typing and DNA similarity analysis implicate *Borrelia valaisiana*-related isolated in Japan is distinguishable from European *B. valaisiana*. *J Vet Med Sci.* 2013;75(9):1201-7.
- 40) Hamaguchi Y, Shirakawa T, Kuwatsuka Y, Ando S. A Neonatal Case of Japanese Spotted Fever. *Pediatr Infect Dis J.* 2013; 32(11): 1286-7.
- 41) Matsutani M, Ogawa M, Takaoka N, Hanaoka N, Toh H, Yamashita A, Oshima K, Hirakawa H, Kuhara S, Suzuki H, Hattori M, Kishimoto T, Ando S, Azuma Y, Shirai M. Complete genome DNA sequence of the East Asian spotted fever disease agent, *Rickettsia japonica*. *PLoS One.* 2013;8(9): e71861.
- 42) Takano A, Fujita H, Kadosaka T, Takahashi M, Yamauchi T, Ishiguro F, Takada N, Yano Y, Oikawa Y, Honda T, Gokuden M, Tsunoda T, Turumi M, Ando S, Andoh M, Sato K, Kawabata H. Construction of a DNA database for ticks collected in Japan: application of molecular identification based on the mitochondrial 16S rDNA gene. *Med Entomol Zool.* 2014;65(1):13-21.
- 43) Gaowa, Yoshikawa Y, Ohashi N, Wu D, Kawamori F, Ikegaya A, Watanabe T, Saitoh K, Takechi D, Murakami Y, Shichi D, Aso K, Ando S. *Anaplasma phagocytophilum* antibodies in humans, Japan, 2010-2011. *Emerg Infect Dis.* 2014;20(3): 508-509.
- 44) Andoh M, Andoh R, Teramoto K, Komiya T, Kaneshima

- T, Takano A, Hayashidani H, Ando S. Survey of *Coxiella burnetii* in ticks collected from dogs in Japan. J Vet Med Sci. 2013;75(8):1115-7.
- 45) Kawabata H, Takano A, Kadosaka T, Fujita H, Nitta Y, Gokuden M, Honda T, Tomida J, Kawamura Y, Masuzawa T, Ishiguro F, Takada N, Yano Y, Andoh M, Ando S, Sato K, Takahashi H, Ohnishi M. Multilocus sequence typing and DNA similarity analysis implicate *Borrelia valaisiana*-related isolated in Japan is distinguishable from European *B. valaisiana*. J Vet Med Sci. 2013;75(9):1201-7.
- 46) Hamaguchi Y, Shirakawa T, Kuwatsuka Y, Ando S. A Neonatal Case of Japanese Spotted Fever. Pediatr Infect Dis J. 2013;32(11):1286-7.
- 47) Matsutani M, Ogawa M, Takaoka N, Hanaoka N, Toh H, Yamashita A, Oshima K, Hirakawa H, Kuhara S, Suzuki H, Hattori M, Kishimoto T, Ando S, Azuma Y, Shirai M. Complete genome DNA sequence of the East Asian spotted fever disease agent, *Rickettsia japonica*. PLoS One. 2013;8(9):e71861.
- 48) Takano A, Fujita H, Kadosaka T, Takahashi M, Yamauchi T, Ishiguro F, Takada N, Yano Y, Oikawa Y, Honda T, Gokuden M, Tsunoda T, Turumi M, Ando S, Andoh M, Sato K, Kawabata H. Construction of a DNA database for ticks collected in Japan: application of molecular identification based on the mitochondrial 16S rDNA gene. Med Entomol Zool. 2014;65(1):13-21.
- 49) Gaowa, Yoshikawa Y, Ohashi N, Wu D, Kawamori F, Ikegaya A, Watanabe T, Saitoh K, Takechi D, Murakami Y, Shichi D, Aso K, Ando S. *Anaplasma phagocytophilum* antibodies in humans, Japan, 2010-2011. Emerg Infect Dis. 2014;20(3):508-9.